



# To Reach the Unreachable: Exploring the Potential of VR Hand Redirection for Upper Limb Rehabilitation

Peixuan Xiong  
The Hong Kong University of Science  
and Technology (Guangzhou)  
Guangzhou, China  
pxiong843@connect.hkust-gz.edu.cn

Yukai Zhang  
The Hong Kong University of Science  
and Technology (Guangzhou)  
Guangzhou, China  
yzhang118@connect.hkust-gz.edu.cn

Nandi Zhang  
University of Calgary  
Calgary, Canada  
nandi.zhang@ucalgary.ca

Shihan Fu  
The Hong Kong University of Science  
and Technology (Guangzhou)  
Guangzhou, China  
sfu663@connect.hkust-gz.edu.cn

Xin Li  
The Third Affiliated Hospital  
Sun Yat-sen University  
Guangzhou, China  
lixin36@mail.sysu.edu.cn

Yadan Zheng  
The Third Affiliated Hospital  
Sun Yat-sen University  
Guangzhou, China  
zhengyad@mail.sysu.edu.cn

Jinni Zhou  
The Hong Kong University of Science  
and Technology (Guangzhou)  
Guangzhou, China  
eejinni@hkust-gz.edu.cn

Xiquan Hu\*  
The Third Affiliated Hospital  
Sun Yat-sen University  
Guangzhou, China  
huxiquan@mail.sysu.edu.cn

Mingming Fan\*  
The Hong Kong University of Science  
and Technology (Guangzhou)  
Guangzhou, China  
The Hong Kong University of Science  
and Technology  
Hong Kong SAR, China  
mingmingfan@ust.hk

## ABSTRACT

Rehabilitation therapies are widely employed to assist people with motor impairments in regaining control over their affected body parts. Nevertheless, factors such as fatigue and low self-efficacy can hinder patient compliance during extensive rehabilitation processes. Utilizing hand redirection in virtual reality (VR) enables patients to accomplish seemingly more challenging tasks, thereby bolstering their motivation and confidence. While previous research has investigated user experience and hand redirection among able-bodied people, its effects on motor-impaired people remain unexplored. In this paper, we present a VR rehabilitation application that harnesses hand redirection. Through a user study and semi-structured interviews, we examine the impact of hand redirection on the rehabilitation experiences of people with motor impairments and its potential to enhance their motivation for upper limb rehabilitation. Our findings suggest that patients are not sensitive to hand movement inconsistency, and the majority express interest in incorporating hand redirection into future long-term VR rehabilitation programs.

\*corresponding author

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](mailto:permissions@acm.org).

CHI '24, May 11–16, 2024, Honolulu, HI, USA

© 2024 Copyright held by the owner/author(s). Publication rights licensed to ACM.

ACM ISBN 979-8-4007-0330-0/24/05

<https://doi.org/10.1145/3613904.3642912>

## CCS CONCEPTS

• **Human-centered computing** → **Empirical studies in accessibility**; *Interaction paradigms*; *Virtual reality*.

## KEYWORDS


Motor impairments, Upper limb rehabilitation, Virtual hand redirection

### ACM Reference Format:

Peixuan Xiong, Yukai Zhang, Nandi Zhang, Shihan Fu, Xin Li, Yadan Zheng, Jinni Zhou, Xiquan Hu, and Mingming Fan. 2024. To Reach the Unreachable: Exploring the Potential of VR Hand Redirection for Upper Limb Rehabilitation. In *Proceedings of the CHI Conference on Human Factors in Computing Systems (CHI '24)*, May 11–16, 2024, Honolulu, HI, USA. ACM, New York, NY, USA, 11 pages. <https://doi.org/10.1145/3613904.3642912>

## 1 INTRODUCTION

Motor impairments, especially hand and arm mobility impairments, are common symptoms among people who live with cerebral palsy (CP), amyotrophic lateral sclerosis (ALS), spinal cord injury (SCI), and brain stroke. Such motor impairments negatively affect their independent living, including performing activities of daily life [33, 85]. Rehabilitation therapies are commonly adopted to help people with motor impairment regain strength and control over their impaired body parts and may help them fully or partially restore their motor ability [10]. Performing rehabilitation exercises in a hospital is a common step that many people with motor impairments (e.g., stroke survivors) take when following rehabilitation therapies. Although there are some methods that allow patients to exercise for rehabilitation without medical equipment, such as games [28, 29], haptic exercises [56], or performing exercises following the initial



**Figure 1: Overview of the study: (A) The participant wears the BURT and the VR headset. BURT helps participants cancel out the burden of gravity on the affected arm. (B) Warm-up for VR Rehabilitation scene with text hint: “Please adjust your hands and body to a comfortable position” (C) The participant performs reaching in VR, corresponding to depth; (D) The participant performs shoulder flexion in VR, corresponding to vertical; (E) The participant performs horizontal abduction in VR, corresponding to horizontal.**

demonstrations from physiotherapists [45]. These methods still cause frustration, tediousness, and lack of motivation [62] because the patients can not perceive their improvement directly. People with motor impairment receiving more physiotherapy treatment recover faster [30, 63]. The majority of rehabilitation hospitals have widely adopted constraint-induced movement therapy (CIMT) for patients, which requires intensive, repetitive daily activities of the affected side while constraining the use of the less-affected side. This approach has demonstrated effective therapeutic outcomes [44, 60]. However, the high intensity of this treatment - demanding more than six hours of daily exercise, five days a week, for a duration exceeding two weeks - also leads to fatigue and reduced compliance for patients [39, 52]. On the other hand, while substantial training efforts might not yield noticeable improvements, patients remain highly sensitive to their failures when employing the affected limbs, discouraging them from engaging in the exercise. As a result, it is important to motivate patients during the lengthy rehabilitation process by allowing them to *perceive improvement* to ensure effective rehabilitation.

Recent developments in Virtual Reality (VR) have shown great potential for making rehabilitation more engaging. VR-based rehabilitation allows patients to carry out rehabilitative exercises while immersing themselves in an interactive virtual environment, increasing the level of engagement which can lead to better therapeutic outcomes [6, 38, 49, 57, 82, 86]. A promising technique for

rehabilitation is *hand redirection*, which breaks the exact one-to-one mapping (i.e. completely synchronized movement) between a patient’s physical hands and the corresponding virtual hands in VR by modifying the visual movements of the virtual hands. With hand redirection, the hand movements of patients with motor impairment can be increased such that they can accomplish seemingly more difficult tasks. This has the potential value to increase their extrinsic motivation. However, such increased movement can potentially reduce users’ sense of hand ownership, which is the feeling that the hand belongs to oneself [5]. The sense of hand ownership can be measured by the noticeability of hand redirection. A reduced sense of hand ownership makes hand redirection more noticeable, which may negatively impact rehabilitation as it reminds patients that their good performance results from external help, thereby reducing their motivation [80]. As a result, it is crucial to find an optimal form of hand redirection that can facilitate improved performance in patients while maintaining a strong sense of hand ownership.

Previous studies on hand redirection, however, are primarily conducted with able-bodied users [50, 84], while little is known about how people with motor impairments perceive hand redirection in VR and how different hand redirection techniques might improve their motivation in rehabilitation. Previous works have explored two primary paradigms: Pre-offset (applying offsets at the initiation of a motion[3, 31], Fig.2 B) or scaling offset (gradually throughout the entire hand movement[9, 20, 31, 42, 84], Fig.2 C). In addition to

these techniques, we propose a new hand redirection method that applies offsets gradually as the hand approaches the target.

Overall, our goal was to increase patients' motivation and effort in rehabilitation while retaining the sense of hand ownership. In this paper, we conducted a user study with 11 patients with upper limb impairments at a local hospital to understand its effect on motivation in comparison with no hand redirection and other hand redirection techniques in the literature. Thus, our research question (RQ) is: **How do different hand redirection techniques affect the effort, motivation, and sense of hand ownership of patients in VR-based rehabilitation?** Our findings show that participants hardly noticed hand redirection and they showed more effort and motivation in VR-based rehabilitation with hand redirection. In summary, we made the following contributions:

- We proposed a novel hand redirection technique that retains the sense of hand ownership while improving participant's training effort and motivation.
- We conducted an empirical study with upper-limb-impaired patients through a VR rehabilitation application that harnesses hand redirection.

## 2 BACKGROUND AND RELATED WORK

### 2.1 Rehabilitation for Upper-limb Motor Impairments

Upper-limb motor impairment is a prominent concern due to its global prevalence and disabling effects. Prioritizing recovery empowers patients for independent daily activities, reducing reliance on caregivers and professionals, and enhancing overall quality of life [16].

Various therapeutic approaches have been explored for stroke patients, including robot-assisted training [4, 12, 26], motor training sessions [15], constraint-induced movement therapy (CIMT) [21, 22, 44, 69], and physical therapy [2, 33, 48]. For instance, Colombo et al. [12] introduced motivating robotic devices for upper limb rehabilitation, incorporating gaming elements (challenges, rewards, rules, etc) to enhance patient engagement and emphasizing regular performance evaluation.

Motivation plays a crucial role in driving improved performance in rehabilitation and is a critical determinant of outcomes [79]. Active participation reflects motivation, while passivity indicates a lack thereof [54]. Traditional methods often struggle to maintain this balance, constrained by the need to moderate training difficulty in light of patient frustration levels [73]. In this work, we tackle this issue by leveraging personalized parameters that are tailored to each participant.

### 2.2 VR for Rehabilitation

Rehabilitating motor-impaired patients aims at promoting effective motor learning [40], a task often hindered by the challenges of intensive training and motivation diminished by low self-efficacy. VR offers a solution, proving effective across conditions including cerebral palsy [53, 58, 67], Parkinson's [47, 66, 81], stroke [17, 19, 35, 38, 72, 75], and injuries [55, 65, 78].

Prior studies have explored the effect of VR in rehabilitation of enhancing patient engagement through visual and auditory

feedback on movement [6, 38, 49, 57, 82, 86]. Zimmerli et al. highlighted the pivotal role of interactivity (ie, functional feedback to motor performance) of VR in engaging patients in rehabilitation [86]. Other studies showcase the effectiveness of combining VR with other approaches, such as conventional physical therapy [34, 36, 59, 61, 70], robotic-assisted rehabilitation [11, 67, 83], and exoskeletons [26, 41, 76] in improving motor functions.

VR has been widely explored as a rehabilitation interface to promote patient motivation in different ways. For instance, Dias et al.'s VR minigames, incorporating competitive elements and encouraging sounds, exemplify strategies to enhance motivation and prevent frustration during rehabilitation [14]. While prior VR interventions often focus on creating motivating content, the potential of leveraging VR's customizability to tackle the self-efficacy-difficulty challenge encountered in real-world rehabilitation remains under-explored. This paper aims to address this gap in the literature.

### 2.3 Hand Redirection in VR

Hand redirection techniques in VR have emerged as an illusion-based method to manipulate the mapping of real-to-virtual hand positions, thereby enhancing user experiences and interactions.


An essential consideration in the implementation of hand redirection techniques is to ensure that users maintain a strong sense of ownership in the virtual environment [51]. Prior studies have demonstrated that inducing a body ownership illusion during VR rehabilitation can improve the therapeutic effect over the non-embodied rehabilitation in VR [8, 24, 64]. As such, a key focus in prior research has been on determining the detection thresholds of these offset across various motion axes [20, 32, 51]. While previous research has explored people's ability to perceive hand redirection in VR environments [84], no prior studies have specifically addressed patients with upper limb impairments. Given that individuals in this patient group typically exhibit lower cognitive abilities, we hypothesized that their detection threshold (i.e., just noticeable difference (JND)) of visual-proprioceptive conflict might be higher than that of healthy individuals. To address this, our research aims to systematically evaluate various hand redirection techniques, examining their impact on patient motivation and the preservation of ownership in VR.

## 3 USER STUDY

To understand how different hand redirection techniques affect the motivation, effort, and sense of hand ownership of people with motor impairments during rehabilitation, we conducted an empirical user study at a local hospital with 11 participants. A VR-based rehabilitation exercise application with three different hand redirection techniques (pre-offset, scaling offset, and post-offset) is implemented. During the study, patients with upper limb impairment are instructed to reach targets in VR at various positions using their affected arm with and without hand redirection because reaching is the most common movement in daily activities and serves as the fundamental movement in stroke rehabilitation training [1].

### 3.1 Hand Redirection Techniques

Three different hand redirection techniques were tested in the study: Pre-offset, Scaling offset, and Post-offset, as illustrated in Fig. 2.



**Figure 2: Hand redirection techniques:** (A) No hand redirection; (B) Pre-offset: the virtual hand is displaced a fixed offset from the real hand from the beginning to the end of hand motion; (C) Scaling offset: the virtual hand scales the movement of the real hand by a fixed factor; (D) Post-offset: the virtual hand gradually enlarging the movement when the real hand approaches its maximum reach. In all offset conditions, the parameters are set such that the virtual hand offsets the real hand the same distance when the real hand is at the maximum reach of the user.

**Pre-offset.** The Pre-offset technique offsets the virtual hand from the physical hand by a constant displacement towards the target regardless of the real hand position. By bringing the affected hand closer to the target, the Pre-offset technique makes it easier for patients to succeed in the target-reaching task, potentially increasing their motivation to engage in the exercise.

**Scaling offset.** The Scaling offset technique scales the position of the virtual hand based on the real hand position from an original position. With a scaling factor larger than 1, it enlarges the virtual hand movement, making it easier for patients to reach the target. Similar to the Pre-offset, this might encourage the patient at the cost of lower training efforts.

However, as the task becomes easier, this may come at the cost of reduced training efforts, harming the therapeutic effect [74]. To resolve this problem, we proposed a new hand redirection technique.

**Post-offset.** Unlike Pre-offset and Scaling-offset, the displacement between the virtual and physical hand for Post-offset is only applied when the physical hand approaches its maximum reach. As the physical hand reaches a threshold close to its maximum reach, a gradually enlarging offset is applied, extending the virtual hand

toward the target until the offset reaches a preset constant displacement. By applying the offset only when the hand approaches its limit, the Post-offset has the potential to induce the needed workload for the desirable therapeutic effect while maintaining a high success rate in accomplishing the tasks, keeping patients motivated.

### 3.2 Apparatus

We conducted the user study in the operational therapy room at a local hospital. We used the Barrett Upper-extremity Robotic Trainer (BURT) to support the patient's affected arm in order to relieve them from the burden of gravity so that more patients with low hand mobility could be included in the study. We conducted the study using the Oculus Quest Pro and Oculus Quest 2 headset, powered by an Intel Core i7 CPU and an NVIDIA GeForce RTX 3070ti GPU. The VR-based rehabilitation application was developed with Unity. With hand tracking, participants can see the virtual representation of their real hands in the VR scene.

**Table 1: Participants' demographic information and their symptoms**

ID	age	Affect side	Motor impairments
1	53	Left	Moderate, able to move the arm against gravity (Grade 3 in MMT)
2	56	Left	Moderate, able to move the arm against gravity (Grade 3 in MMT)
3	51	Left	Moderate, able to move the arm against gravity (Grade 3 in MMT)
4	51	Left	Mild, able to move the arm against gravity and resistance (Grade 4 in MMT)
5	66	Left	Moderate, able to move the arm against gravity (Grade 3 in MMT)
6	66	Right	Mild, able to move the arm against gravity and resistance (Grade 4 in MMT)
7	52	Right	Moderate, able to move the arm against gravity (Grade 3 in MMT)
8	60	Right	Mild, able to move the arm against gravity and resistance (Grade 4 in MMT)
9	22	Left	Severe, able to move the arm without gravity (Grade 1 in MMT)
10	29	Left	Severe, able to move the arm without gravity (Grade 1 in MMT)
11	50	Left	Severe, able to move the arm without gravity (Grade 1 in MMT)

### 3.3 Participants

We recruited 14 patients from the local hospital, and three of them discontinued their participation during the study because they were uncomfortable with VR devices (Reported dizzy or tired). Consequently, our final sample consisted of 11 patients with upper limb motor impairments, with an average age of 50.5. Our recruitment criteria were based on the extensor and flexor muscles achieving grades 1-4 based on manual muscle testing (MMT) [13] on hands. Participants in these states have a small amount of athletic ability on their hands or lack of endurance. Before official recruitment, we assessed their mental state using the Minimum Mental State Examination (MMSE) [43], and all participants demonstrated normal cognitive abilities to express emotion and perform simple calculations and got grades higher than 22 in MMSE. None of the participants had prior experience with VR. Approval was obtained from the ethical committees of our home institution and the hospital. All participants gave us written informed consent and received a \$10 shopping card for compensation. Detailed demographics are shown in Table. 1.

### 3.4 Design

We designed a within-subject study with the following independent variables: 1) **Hand redirection technique**. it has five levels: No hand redirection, No hand redirection with reduced difficulty, Pre-offset, Scaling offset, and Post-offset. Three different hand redirection techniques are investigated in the study. The parameters for the hand redirection techniques are configured such that the displacement between the virtual and the physical hands is the same across all offset conditions when the users reach maximum reach in the designated direction. *No hand redirection* is used as a baseline. However, hand redirection effectively reduces the task difficulty by reducing the required travel distance of the user's hand. To understand the effect of hand redirection with tasks of the same difficulty, we included a second baseline - *no hand redirection with reduced difficulty*, in which the target is set closer to the user's hand such that its task difficulty is the same as the three offset conditions. 2) **Hand movement direction**. It has three levels: horizontal, vertical, and depth. The direction in which the target shows up for the participant to reach.

Our dependent variables include 1) **Training efforts**. We used the average hand travel distance in each trial to measure it. We

assume that more effect is spent on training when the affected hand reaches further toward the target direction. To account for differences in each participant's ability, we normalized the hand travel distance by dividing it by its maximum reach distance recorded during the preparation stage.; 2) **Sense of hand ownership**. This is measured by a questionnaire in which participants reported their level of avatar embodiment on hands [27] on a 7-point Likert scale.

### 3.5 Procedure

**Preparation and warm-up.** Before the study, participants were told that the study investigated the efficacy of rehabilitation in VR. They are not informed of the real research target. After signing the informed consent form, participants were assisted in donning the equipment and positioned in front of the BURT Rehabilitation Robot. Their affected hand was securely affixed to the apparatus, as shown in (Fig. 1 A). Participants found themselves in a laboratory setting inside the VR environment with an empty table before them. They could see their affected hand resting on the table. Participants were informed that their mission was to reach the red ball, and if its color changed to black, they would consider it a success and should return their hand to the initial position. In cases where participants couldn't accomplish the mission, they were instructed to return their hands to the initial position and wait for another refresh. Each participant was given one minute to explore the virtual environment to get familiar with hand movement and the task. During this warm-up section, no hand redirection techniques were applied.

**Rehabilitation exercise.** Participants were instructed to adjust their hands and body to a comfortable position (Fig. 1 B). Following this, participants were directed to perform specific movements involving their hands' reaching, flexion, and horizontal abduction to the fullest extent possible along the x, y, and z axes, corresponding to depth, vertical, and horizontal, respectively (Fig. 1 C, D, E). These movements are commonly used for rehabilitation [18]. The maximum reach distances in these three directions were stored. Moving forward, a ball was generated relative to the patient's hand's initial position. This target was positioned along the x, y, and z axes, with distances set at 1, 1.25, and 1.5 times the patient's maximum achievable range of motion (calculated from the Max Position to the Initial Position). For each of the three distances, participants completed one action under 1.25 times offset on the x-axis, then three reaching



actions under 1.5 times offset. The target refreshes six times on each axis if it is a baseline condition. Each patient experienced five conditions in a randomized order for counterbalancing. The patient will rest and take up the VR headset between each condition to complete the questionnaire and answer the Likert scale questions. This exercise took around 30 minutes.

**Semi-structured interview.** We conducted a semi-structured interview with nine participants. Two of the 11 participants refused the interview due to personal reasons. During this interview session, we provided participants with information about our genuine research objectives and demonstrated the practical application of hand redirection within the VR environment. We engaged them in discussions about the effects of their efforts, motivation, and sense of hand ownership in VR-based rehabilitation. Each interview session lasted approximately 20 minutes for each participant.

### 3.6 Data Analysis

For each participant we obtained 90 trials (3 target distance $\times$ 2 coefficient $\times$ 3 direction $\times$ 5 technique), and 11 participants achieved 990 trials in total. We pre-processed the data by eliminating invalid trails caused by inaccurate hand-tracking (The hands are not detected) and participants could not move their hands due to severe motor impairments in the designated direction (Sum up to 60 trials). The Shapiro-Wilk normality test was executed upon eliminating the invalid trails, revealing that the data from questionnaires did not follow a normal distribution. Still, data from travel distances of hands followed a normal distribution. Consequently, the Friedman test [25] was conducted to assess the difference between data in questionnaires and two-way repeated-measures ANOVA was conducted to assess the difference between data in travel distances.

## 4 RESULTS AND FINDINGS

In this section, we will combine the results of data from participants and open coding interviews to investigate how hand redirection affects people with motor impairments in effort and performance, motivation, and ownership.

### 4.1 Effort and Performance

Questionnaire results show no significant differences between different hand redirection techniques on self-reported performance ( $p = 0.818$ ) and effort ( $p = 0.115$ ).

Figure 3 shows the effect of different hand direction techniques on effort measured by the hand travel distance. Two-way repeated-measures ANOVA results show a significant difference in the participants' effort to reach the target between different hand redirection techniques ( $p < .001$ ) and movement directions ( $p < .001$ ). Among all conditions, participants spent significantly higher efforts in offset conditions (*Pre-offset*, *Scaling offset*, and *Post offsets*) comparing to the baseline (*No hand redirection with Reduced difficulty*), which means hand redirection can increase patients' effort in reaching tasks during rehabilitation exercise when the task difficulty is the same. Among the three techniques, *Post-offset* yields the highest increase in effort.

### 4.2 Motivation

During the interview, almost all participants thought that hand redirection could enhance their motivation, as it helped them complete the task. (P4: "I feel frustrated when I can not reach the target, so I want to be helped." P5: "It is better to deploy hand redirection, which can improve the range of movement. If I can not reach the target, I will give up. This can improve my confidence because my confidence comes from completing tasks. If I perform worse, he will lose confidence and motivation. For long-term rehabilitation, this will also improve my motivation.") Only one participant (P10) rejected it as he preferred to see his truthful capability in the virtual scene during exercises.

When asked whether the task difficulty should be reduced by placing targets closer to the affected hand so that they can reach more targets, most participants objected. Some participants believe that closer targets will make the exercise boring (P4: "It is boring when the target is too near to the hand.", P5: "It seems too easy to reach, and makes it boring, losing interest."). Other participants worried that easier tasks may hinder the therapeutic effect. (P6: "The task should be more difficult because I exercise here. Too easy may bring no effect on rehabilitation")


### 4.3 Sense of Hand Ownership

Results from interviews and questionnaires show that participants hardly notice hand redirection, which implies a high sense of hand ownership. RM-ANOVA test shows no significant differences between the five conditions on self-reported hand movement consistency ( $p = 0.39$ ), hand location consistency ( $p = 0.745$ ), and hand ownership ( $p = 0.123$ ). During the interview, when asked whether they noticed there was displacement between the real hand and the virtual hand, only two participants said they did. After revealing our true research objectives and demonstrating different hand redirection techniques, two more participants (P4, P6) said that they noticed the displacement but they did not report it as they were unable to articulate their observation so they chose to disregard them (P6: "There are differences in my hands, But I can not tell what exactly the differences are. The hands in the VR seem less flexible.").

### 4.4 Preference on Hand Redirection Techniques

Interview results show most participants prefer to use the *Post-offset* for future VR-based rehabilitation exercises. During the interview, participants were asked to choose a technique for future rehabilitation in VR. While two participants have no preference (P4, P8: "All methods are great because I can reach more targets. I can accept hand does not belong to me in VR rehabilitation"), other participants had the following preferences:

**Post-offset.** Most participants prefer the *Post-offset* for two different reasons: 1) *Post-offsets* could enhance their endurance (P3: "I want to be helped at the end when I can not reach the target." The pleasure lies in the accomplishment achieved through diligent effort", P6: "In *Post-offset* condition, the hand is reaching to target slowly. It'll build up my endurance.") 2) It encourages the participant to try harder. (P5: "Help me at the end is a better technique because there is an expected value that I will not easily give up. As for the other two redirection methods, the task seems easier when *offset* appears since



**Figure 3: The training effort in three movement directions (depth, vertical, and horizontal) with five different hand redirection conditions, including three hand redirection techniques (Pre-offset, Scaling offset, and Post-offset) and two baseline conditions (no hand redirection w/o reduced difficulty).**

the beginning of each trial. Then, I will not try harder and may give up.)

**Scaling offset.** P7 preferred Scaling offset as it is more entertaining (P7: "The Post-offset is similar to Scaling offset, so it is hard to say which is better. But I will choose Scaling offset because it is much more exciting than other techniques.").

**Pre-offset.** P9 preferred Pre-offset due to its low noticeability. However, other participants suggested that once it is noticed, the Pre-offset felt unrealistic (P6: "It is disproportionate." P7: "It is not realistic. Moreover, it is much more excessive than Scaling offset.").

**Additional comments on the systems.** The system should incorporate haptic feedback. (P5: "The environment is great and real, but the appearance of the hands is not real enough. The haptic of hands is also lacking. When I can not feel haptic from my disabled hand, I try another hand, but I still can not feel anything.") We also find this phenomenon in other participants. They will try their other hand to seek haptic feedback. This participant also gave us feedback compared with other devices in daily rehabilitation in the hospital. I felt as if addicted to it. When using this system, it is not easy to make me feel tired. This is far more interesting than other devices I experienced in daily rehabilitation" Young people may be less likely to accept hand redirection. (P6: "The young patients may not easily accept this method. They are sensitive to the help from the

machine.") But the P9 disagrees with that. (P9: "I played so many computer games so I can notice the difference between this VR training and other training. I am in a game. I do not care if I can push my hand to limit this way.")

## 5 DISCUSSION AND FUTURE WORK

We investigated how hand redirection affects rehabilitation for people with upper-limb motor impairments. In summary, our findings show that hand redirection improves the training effort without losing the sense of hand ownership for people with motor impairments. Furthermore, almost all participants expressed a strong inclination toward incorporating hand redirection. They believed it could enhance their confidence in the rehabilitation process by completing more missions, and they will be more motivated [77]. Among all hand redirection conditions, the proposed Post-offset enhanced participants' effort and performance and kept more ownership, while participants felt it to be more motivating. Below we discussed our findings.

### 5.1 Training Effort

**Visually more difficult tasks lead to higher efforts.** Participants reach farther with hand redirection than the baseline condition with reduced difficulty. This might be the result of the smaller visual

target distance in the baseline with reduced difficulty which makes participants subconsciously put in less effort.

**Reduced task difficulty resulting from hand redirection can lead to lower efforts.** It is more difficult for participants to reach the target in *no hand redirection* condition compared to offset conditions as hand redirection effectively reduces the required travel distance of the hand for reaching the target. Our result shows that participants spend significantly less effort in *Pre-offset* and *Scaling offset*. *Post-offset* also exhibits less training effort compared to *no hand redirection* in two directions though there is no statistical significance. These results show that reduced task difficulty results from hand redirection can lead to lower efforts.

**Post-offset outperforms Pre-offsets and Scaling offsets in retaining training effort.** Participants spent significantly more effort in rehabilitation exercises with *Post-offset* compared to other offset conditions. The rationale behind this was revealed in the interview in which participants suggested that *Post-offset* only helps them at the end when they cannot reach the target, which can be helpful in building up their endurance. These results show that *Post-offset* has the potential to reduce task difficulty while avoiding slacking, which could decrease patients' motor output and decrease recovery [7].

**Participants reach farther in the depth direction.** This is because the participant can lean forward their body to compensate for their impairment when reaching the target. Compensation is common because people with motor impairment, as it is necessary for completing most activities of daily living [37].

## 5.2 Motivation

**Participants believe hand redirection can enhance their motivation in these rehabilitation tasks** According to our results from semi-structured interviews, almost all participants in this study reported they would like to use hand redirection in future VR rehabilitation. They believe it could enhance their motivation and confidence in the long term due to more tasks they can achieve, which is also reflected by the results of the efforts. In future work, we can conduct a longitudinal rehabilitation study to evaluate how hand redirection motivates patients in the long term. We can also combine the evaluation results in the hospital, and compare the training effect between VR-based rehabilitation with hand redirection and traditional training methods.

**Participants object to reducing task difficulty in an obvious manner** Although participants consider hand redirection techniques (which reduce task difficulty) helpful, most participants ( $N = 10$ ) objected to reducing task difficulty by simply placing the target closer as they think it makes the exercise less challenging and may harm the therapeutic effect. This suggests that patients may prefer tasks that are visually more challenging during rehabilitation exercises. This may be explained by the comfort zone learning model, which theorizes that learners learn better when tasks are more difficult and outside of their comfort zone [46]. Future work could investigate how visual difficulty (visually challenging or complex tasks [68]) affects the motivation and effectiveness of rehabilitation.

## 5.3 Virtual Hand Ownership

**Participants hardly notice hand redirection.** Most participants could not notice the difference between different conditions and gave full points for a sense of hand ownership in questionnaires. This finding was corroborated by the results of semi-structured interviews, which show most participants are not aware of the disparities between the virtual hand and the real hand. In future work, we may use quantitative methods such as EEG [23] or EMG [71] to investigate how hand redirection affects patients' sense of hand ownership and help explore the optimal hand redirection threshold in people with motor impairments.

## 6 LIMITATION

While our results suggest promise of VR hand redirection for rehabilitation, this work is limited in the following aspects. First, medical evaluation is not included in the study. Due to limited time and resources, each participant only experienced the VR-based rehabilitation application for less than one hour which is too short for any noticeable therapeutic outcomes. Future work can investigate the long-term effect of hand redirection on therapeutic outcomes. Second, our results may not apply to patients with mild symptoms as the study was conducted with hospitalized patients with upper limb muscle strength levels between 1 to 4, while people with higher muscle strength might exhibit less sense of hand ownership due to better proprioception. Third, the warm-up section lasts for only 1 minute. This may cause the participants to remain unfamiliar with VR and hand redirection and make it harder to find virtual hand displacements. Fourth, VR equipment and app interface can also impact patient motivation and embodiment in VR due to insufficiently realistic environments and unfamiliar operation methods. This could be investigated through questionnaires in future work. Finally, although we have selected reasonable parameters (e.g. target distances, displacement for hand redirection) for VR-based rehabilitation exercises with hand redirection by testing with abled-bodied people, these parameters are not optimal. Future work can investigate the optimal value of displacement for hand redirection or target distances in VR-based rehabilitation in general.

## 7 CONCLUSION

Hand redirection holds immense potential in improving the motivation of people with upper limb motor impairment during VR-based rehabilitation exercises. While prior research in hand redirection mostly focuses on the experience of hand redirection for abled-bodied people, this study examined the effect of various hand redirection techniques on the motivation, effort, and sense of hand ownership of people with upper limb motor impairment during VR-based rehabilitation exercises. We proposed a novel hand redirection technique - *Post offset* for upper limb rehabilitation. A user study was conducted at a local hospital with 11 participants with upper limb motor impairment. Our findings suggest that while people with upper limb motor impairment can hardly notice hand redirection, hand redirection improves their motivation and effort during VR rehabilitation exercises. Participants show the highest effort with *Post-offset* among all hand redirection techniques investigated in this study. Most participants express interest in incorporating hand redirection into future long-term VR rehabilitation programs.



## ACKNOWLEDGMENTS

This work is partially supported by 1) 2024 Guangzhou Science and Technology Program City-University Joint Funding Project (PI: Mingming Fan); 2) 2023 Guangzhou Science and Technology Program City-University Joint Funding Project (No. 2023A03J0001) ; 3) Guangdong Provincial Key Lab of Integrated Communication, Sensing and Computation for Ubiquitous Internet of Things (No.2023B1212010007).

## REFERENCES

- [1] Margit Alt Murphy and Charlotte K Häger. 2015. Kinematic analysis of the upper extremity after stroke—how far have we reached and what have we grasped? *Physical Therapy Reviews* 20, 3 (2015), 137–155.
- [2] Saba Anwer, Asim Waris, Syed Omer Gilani, Javaid Iqbal, Nusratnaaz Shaikh, Amit N. Pujari, and Imran Khan Niazi. 2022. Rehabilitation of Upper Limb Motor Impairment in Stroke: A Narrative Review on the Prevalence, Risk Factors, and Economic Statistics of Stroke and State of the Art Therapies. *Healthcare* 10, 2 (2022). <https://doi.org/10.3390/healthcare10020190>
- [3] Brett Benda, Shaghayegh Esmaili, and Eric D. Ragan. 2020. Determining Detection Thresholds for Fixed Positional Offsets for Virtual Hand Remapping in Virtual Reality. In *2020 IEEE International Symposium on Mixed and Augmented Reality (ISMAR)*. 269–278. <https://doi.org/10.1109/ISMAR50242.2020.00050>
- [4] Amy A. Blank, James A. French, Ali Utku Pehlivan, and Marcia K. O'Malley. 2014. Current Trends in Robot-Assisted Upper-Limb Stroke Rehabilitation: Promoting Patient Engagement in Therapy. *Current Physical Medicine and Rehabilitation Reports* 2, 3 (2014), 184–195. <https://doi.org/10.1007/s40141-014-0056-z>
- [5] Olaf Blanke. 2012. Multisensory brain mechanisms of bodily self-consciousness. *Nature Reviews Neuroscience* 13, 8 (2012), 556–571.
- [6] Desiderio Cano Porras, Petra Siemonsma, Rivka Inzelberg, Gabriel Zeilig, and Meir Plotnik. 2018. Advantages of virtual reality in the rehabilitation of balance and gait: Systematic review. *Neurology* 90, 22 (May 2018), 1017–1025.
- [7] Maura Casadio and Vittorio Sanguineti. 2012. Learning, retention, and slacking: a model of the dynamics of recovery in robot therapy. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* 20, 3 (2012), 286–296.
- [8] Kuan Cha, Jinying Wang, Yan Li, Longbin Shen, Zhuoming Chen, and Jinyi Long. 2021. A novel upper-limb tracking system in a virtual environment for stroke rehabilitation. *Journal of NeuroEngineering and Rehabilitation* 18, 1 (2021), 1–13.
- [9] Lung-Pan Cheng, Eyal Ofek, Christian Holz, Hrvoje Benko, and Andrew Wilson. 2017. Sparse Haptic Proxy: Touch Feedback in Virtual Environments Using a General Passive Prop. 3718–3728. <https://doi.org/10.1145/3025453.3025753>
- [10] Yaqi Chu, Xingang Zhao, Yijun Zou, Weiliang Xu, and Yiwen Zhao. 2018. Robot-Assisted Rehabilitation System Based on SSVEP Brain-Computer Interface for Upper Extremity. In *2018 IEEE International Conference on Robotics and Biomimetics (ROBIO)*. 1098–1103. <https://doi.org/10.1109/ROBIO.2018.8664812>
- [11] William E Clark, Manoj Sivan, and Rory J O'Connor. 2019. Evaluating the use of robotic and virtual reality rehabilitation technologies to improve function in stroke survivors: A narrative review. *Journal of Rehabilitation and Assistive Technologies Engineering* 6 (2019), 2055668319863557. <https://doi.org/10.1177/2055668319863557> arXiv:<https://doi.org/10.1177/2055668319863557> PMID: 31763052.
- [12] Roberto Colombo, Fabrizio Pisano, Alessandra Mazzone, Carmen Delconte, Silvestro Micera, M Chiara Carrozza, Paolo Dario, and Giuseppe Minuco. 2007. Design strategies to improve patient motivation during robot-aided rehabilitation. *Journal of NeuroEngineering and Rehabilitation* 4, 3 (2007). <https://doi.org/10.1186/1743-0003-4-3>
- [13] Scott C Cuthbert and George J Goodhead Jr. 2007. On the reliability and validity of manual muscle testing: a literature review. *Chiropractic & osteopathy* 15, 1 (2007), 4.
- [14] Paulo Dias, Ricardo Silva, Paula Amorim, Jorge Lains, Eulália Roque, Inês Seródio, Fátima Pereira, and Beatriz Sousa Santos. 2019. Using Virtual Reality to Increase Motivation in Poststroke Rehabilitation. *IEEE Computer Graphics and Applications* 39, 1 (2019), 64–70. <https://doi.org/10.1109/MCG.2018.2875630>
- [15] Volker Dietz, Roland Müller, and Gery Colombo. 2002. Locomotor activity in spinal man: significance of afferent input from joint and load receptors. *Brain* 125, 12 (12 2002), 2626–2634. <https://doi.org/10.1093/brain/awf273> arXiv:<https://academic.oup.com/brain/article-pdf/125/12/2626/1115318/awf273.pdf>
- [16] Marcel Dijkers. 1997. Quality of life after spinal cord injury: a meta analysis of the effects of disablement components. *Spinal Cord* 35 (1997), 829–840. <https://doi.org/10.1038/sj.sc.3100571>
- [17] Jared Duval, Rutul Thakkar, Delong Du, Kassandra Chin, Sherry Luo, Aviv Elor, Magy Seif El-Nasr, and Michael John. 2022. Designing Spellcasters from Clinician Perspectives: A Customizable Gesture-Based Immersive Virtual Reality Game for Stroke Rehabilitation. *ACM Trans. Access. Comput.* 15, 3, Article 26 (aug 2022), 25 pages. <https://doi.org/10.1145/3530820>
- [18] Michael D Ellis, Carolina Carmona, Justin Drogos, and Julius PA Dewald. 2018. Progressive abduction loading therapy with horizontal-plane viscous resistance targeting weakness and flexion synergy to treat upper limb function in chronic hemiparetic stroke: a randomized clinical trial. *Frontiers in neurology* 9 (2018), 71.
- [19] Aviv Elor, Mircea Teodorescu, and Sri Kurniawan. 2018. Project Star Catcher: A Novel Immersive Virtual Reality Experience for Upper Limb Rehabilitation. *ACM Trans. Access. Comput.* 11, 4, Article 20 (nov 2018), 25 pages. <https://doi.org/10.1145/3265755>
- [20] Shaghayegh Esmaili, Brett Benda, and Eric D. Ragan. 2020. Detection of Scaled Hand Interactions in Virtual Reality: The Effects of Motion Direction and Task Complexity. In *2020 IEEE Conference on Virtual Reality and 3D User Interfaces (VR)*. 453–462. <https://doi.org/10.1109/VR46266.2020.00066>
- [21] Mohammad Etoom, Mohammad Hawamdeh, Ziad Hawamdeh, Mohammad Alwardat, Laura Giordani, Serenella Bacciu, Claudia Scarpini, and Calogero Foti. 2016. Constraint-induced movement therapy as a rehabilitation intervention for upper extremity in stroke patients: systematic review and meta-analysis. *Int. J. Rehabil. Res.* 39, 3 (Sept. 2016), 197–210.
- [22] S Fabbri, G Casati, and D Bonaiuti. 2014. Is CIMT a rehabilitative practice for everyone? Predictive factors and feasibility. *Eur. J. Phys. Rehabil. Med.* 50, 5 (Oct. 2014), 505–514.
- [23] Martin Feick, Kora P Regitz, Anthony Tang, Tobias Jungbluth, Maurice Rekrut, and Antonio Krüger. 2023. Investigating Noticeable Hand Redirection in Virtual Reality using Physiological and Interaction Data. In *2023 IEEE Conference Virtual Reality and 3D User Interfaces (VR)*. IEEE, 194–204.
- [24] Giulia Fregna, Nicola Schincaglia, Andrea Baroni, Sofia Straudi, and Antonino Casile. 2022. A novel immersive virtual reality environment for the motor rehabilitation of stroke patients: A feasibility study. *Frontiers in Robotics and AI* 9 (2022), 906424.
- [25] Milton Friedman. 1940. A comparison of alternative tests of significance for the problem of m rankings. *The annals of mathematical statistics* 11, 1 (1940), 86–92.
- [26] Antonio Frisoli, Massimo Bergamasco, Maria C Carboncini, and Bruno Rossi. 2009. Robotic assisted rehabilitation in Virtual Reality with the L-EXOS. *Stud. Health Technol. Inform.* 145 (2009), 40–54.
- [27] Mar Gonzalez-Franco and Tabitha C Peck. 2018. Avatar embodiment: towards a standardized questionnaire. *Frontiers in Robotics and AI* 5 (2018), 74.
- [28] Maja Gorsic, Imre Cikajlo, Metka Javh, Nika Goljar, and Domen Novak. 2019. Pilot long-term evaluation of competitive and cooperative exercise games in inpatient stroke rehabilitation. In *2019 IEEE 16th International Conference on Rehabilitation Robotics (ICORR)*. IEEE, 648–653.
- [29] Maja Gorsic and Domen Novak. 2016. Design and pilot evaluation of competitive and cooperative exercise games for arm rehabilitation at home. In *2016 38th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*. IEEE, 4690–4694.
- [30] Glasgow Augmented Physiotherapy Study (GAPS) Group. 2004. Can augmented physiotherapy input enhance recovery of mobility after stroke? A randomized controlled trial. *Clinical rehabilitation* 18, 5 (2004), 529–537.
- [31] Dustin T. Han, Mohamed Suhail, and Eric D. Ragan. 2018. Evaluating Remapped Physical Reach for Hand Interactions with Passive Haptics in Virtual Reality. *IEEE Transactions on Visualization and Computer Graphics* 24, 4 (2018), 1467–1476. <https://doi.org/10.1109/TVCG.2018.2794659>
- [32] Judith Hartfill, Jenny Gabel, Lucie Kruse, Susanne Schmidt, Kevin Riebandt, Simone Kühn, and Frank Steinicke. 2021. Analysis of Detection Thresholds for Hand Redirection during Mid-Air Interactions in Virtual Reality. In *Proceedings of the 27th ACM Symposium on Virtual Reality Software and Technology (Osaka, Japan) (VRST '21)*. Association for Computing Machinery, New York, NY, USA, Article 35, 10 pages. <https://doi.org/10.1145/3489849.3489866>
- [33] Samar M Hatem, Geoffroy Saussez, Margaux Della Faille, Vincent Prist, Xue Zhang, Delphine Dispa, and Yannick Bleyenheuff. 2016. Rehabilitation of motor function after stroke: a multiple systematic review focused on techniques to stimulate upper extremity recovery. *Frontiers in human neuroscience* 10 (2016), 442.
- [34] Hsiu-Yun Hsu, Li-Chieh Kuo, Yu-Ching Lin, Fong-Chin Su, Tai-Hua Yang, and Che-Wei Lin. 2022. Effects of a virtual reality-based mirror therapy program on improving sensorimotor function of hands in chronic stroke patients: A randomized controlled trial. *Neurorehabil. Neural Repair* 36, 6 (June 2022), 335–345.
- [35] Qianqian Huang, Wei Wu, Xiaolong Chen, Bo Wu, Longqiang Wu, Xiaoli Huang, Songhe Jiang, and Lejian Huang. 2019. Evaluating the effect and mechanism of upper limb motor function recovery induced by immersive virtual-reality-based rehabilitation for subacute stroke subjects: study protocol for a randomized controlled trial. *Trials* 20, 1 (Feb. 2019), 104.
- [36] Ng Zhilong Jerome-Christian, B. S. Rajaratnam, and Tim Xu Tian. 2012. Can Virtual Reality Integrated within Conventional Rehabilitation Enhance Balance Recovery after Stroke?. In *Proceedings of the 6th International Conference on Rehabilitation Engineering & Assistive Technology (Tampines, Singapore) (i-CREATE '12)*. Singapore Therapeutic, Assistive & Rehabilitative Technologies (START)

- Centre, Midview City, SGP, Article 34, 4 pages.
- [37] Theresa A Jones. 2017. Motor compensation and its effects on neural reorganization after stroke. *Nature Reviews Neuroscience* 18, 5 (2017), 267–280.
  - [38] Younbo Jung, Shih-Ching Yeh, and Jill Campbell Stewart. 2006. Tailoring virtual reality technology for stroke rehabilitation: A human factors design. *Conference on Human Factors in Computing Systems - Proceedings*, 929–934. <https://doi.org/10.1145/1125451.1125631>
  - [39] Jeong-Hui Kim and Moon-Young Chang. 2018. Effects of modified constraint-induced movement therapy on upper extremity function and occupational performance of stroke patients. *Journal of Physical Therapy Science* 30, 8 (2018), 1092–1094.
  - [40] Pawel Kiper, Andrzej Szczudlik, Michela Agostini, Jozef Opara, Roman Nowobilski, Laura Ventura, Paolo Tonin, and Andrea Turolla. 2018. Virtual reality for upper limb rehabilitation in subacute and chronic stroke: A randomized controlled trial. *Arch. Phys. Med. Rehabil.* 99, 5 (May 2018), 834–842.e4.
  - [41] Verena Klamroth-Marganska, Javier Blanco, Katrin Campen, Armin Curt, Volker Dietz, Thierry Ettlin, Morena Felder, Bernd Fellinghauer, Marco Guidali, Anja Kollmar, Andreas Luft, Tobias Nef, Corina Schuster-Amft, Werner Stahel, and Robert Riener. 2014. Three-dimensional, task-specific robot therapy of the arm after stroke: a multicentre, parallel-group randomised trial. *Lancet Neurol.* 13, 2 (Feb. 2014), 159–166.
  - [42] Luv Kohli. 2013. *Redirected touching*. Ph. D. Dissertation. The University of North Carolina at Chapel Hill.
  - [43] Lenore Kurlowicz and Meredith Wallace. 1999. The mini-mental state examination (MMSE). , 8–9 pages.
  - [44] Gert Kwakkel, Marc Rietberg, and Erwin van Wegen. 2007. Constraint-induced movement therapy improves upper extremity motor function after stroke. *Australian Journal of Physiotherapy* 53, 2 (2007), 132.
  - [45] Agnes WK Lam, Ahmed HajYasien, and Dana Kulic. 2014. Improving rehabilitation exercise performance through visual guidance. In *2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*. IEEE, 1735–1738.
  - [46] Sarah I Leberman and Andrew J Martin. 2002. Does pushing comfort zones produce peak learning experiences? *Journal of Outdoor and Environmental Education* 7, 1 (2002), 10–19.
  - [47] Cheng Lei, Kejimu Sunzi, Fengling Dai, Xiaoqin Liu, Yanfen Wang, Baolu Zhang, Lin He, and Mei Ju. 2019. Effects of virtual reality rehabilitation training on gait and balance in patients with Parkinson's disease: A systematic review. *PLoS One* 14, 11 (Nov. 2019), e0224819.
  - [48] Mindy F Levin and Marika Demers. 2021. Motor learning in neurological rehabilitation. *Disabil. Rehabil.* 43, 24 (Dec. 2021), 3445–3453.
  - [49] Gwyn N Lewis and Juliet A Rosie. 2012. Virtual reality games for movement rehabilitation in neurological conditions: how do we meet the needs and expectations of the users? *Disabil. Rehabil.* 34, 22 (April 2012), 1880–1886.
  - [50] Zhipeng Li, Yu Jiang, Yihao Zhu, Ruijia Chen, Ruolin Wang, Yuntao Wang, Yukang Yan, and Yuanchun Shi. 2022. Modeling the Noticeability of User-Avatar Movement Inconsistency for Sense of Body Ownership Intervention. *Proceedings of the ACM on Interactive, Mobile, Wearable and Ubiquitous Technologies* 6, 2 (2022), 1–26.
  - [51] Zhipeng Li, Yu Jiang, Yihao Zhu, Ruijia Chen, Ruolin Wang, Yuntao Wang, Yukang Yan, and Yuanchun Shi. 2022. Modeling the Noticeability of User-Avatar Movement Inconsistency for Sense of Body Ownership Intervention. *Proc. ACM Interact. Mob. Wearable Ubiquitous Technol.* 6, 2, Article 64 (jul 2022), 26 pages. <https://doi.org/10.1145/3534590>
  - [52] K-C Lin, C-Y Wu, T-H Wei, Chang Gung, C-Y Lee, and J-S Liu. 2007. Effects of modified constraint-induced movement therapy on reach-to-grasp movements and functional performance after chronic stroke: a randomized controlled study. *Clinical rehabilitation* 21, 12 (2007), 1075–1086.
  - [53] Wei Liu, Yuanyan Hu, Junfeng Li, and Chang Jindong. 2022. Effect of Virtual Reality on Balance Function in Children With Cerebral Palsy: A Systematic Review and Meta-analysis. *Frontiers in Public Health* 10 (04 2022), 865474. <https://doi.org/10.3389/fpubh.2022.865474>
  - [54] Niall Maclean, Pandora Pound, Charles Wolfe, and Anthony Rudd. 2002. The Concept of Patient Motivation. *Stroke* 33, 2 (2002), 444–448. <https://doi.org/10.1161/hs0202.102367> arXiv:<https://www.ahajournals.org/doi/pdf/10.1161/hs0202.102367>
  - [55] Giuseppa Maresca, Maria Grazia Maggio, Antonio Buda, Gianluca La Rosa, Alfredo Manuli, Placido Bramanti, Rosaria De Luca, and Rocco Salvatore Calabrò. 2018. A novel use of virtual reality in the treatment of cognitive and motor deficit in spinal cord injury: A case report. *Medicine* 97, 50 (2018).
  - [56] Mostefa Masmoudi, Nadia Zenati, Samir Benbelkacem, Zineb Hadjadj, Oualid Djekoune, Mohamed Amine Guerrouji, and Youssa Izountar. 2021. Low-cost haptic glove for grasp precision improvement in Virtual Reality-Based Post-Stroke Hand Rehabilitation. In *2021 International Conference on Artificial Intelligence for Cyber Security Systems and Privacy (AI-CSP)*. IEEE, 1–3. <https://doi.org/10.1109/AI-CSP52968.2021.9671087>
  - [57] Alma S Merians, David Jack, Rares Boian, Marilyn Tremaine, Grigore C Burdea, Sergei V Adamovich, Michael Recce, and Howard Poizner. 2002. Virtual reality-augmented rehabilitation for patients following stroke. *Phys. Ther.* 82, 9 (Sept. 2002), 898–915.
  - [58] Burcu Metin Ökmen, Meryem Doğan Aslan, Güldal Funda Nakipoğlu Yüzer, and Neşe Özgürin. 2019. Effect of virtual reality therapy on functional development in children with cerebral palsy: A single-blind, prospective, randomized-controlled study. *Turk. J. Phys. Med. Rehabil.* 65, 4 (June 2019), 371–378.
  - [59] Manasa S Nair, Vivek N Kulkarni, and Ashok K Shyam. 2022. Combined effect of virtual Reality Training (VRT) and conventional therapy on sitting balance in patients with spinal cord injury (SCI): Randomized control trial. *Neurol. India* 70, Supplement (Sept. 2022), S245–S250.
  - [60] Rinske Nijland, Gert Kwakkel, Japie Bakers, and Erwin van Wegen. 2011. Constraint-induced movement therapy for the upper paretic limb in acute or sub-acute stroke: a systematic review. *International Journal of Stroke* 6, 5 (2011), 425–433.
  - [61] Young-Bin Oh, Gi-Wook Kim, Kap-Soo Han, Yu Hui Won, Sung-Hee Park, Jeong-Hwan Seo, and Myoung-Hwan Ko. 2019. Efficacy of virtual reality combined with real instrument training for patients with stroke: A randomized controlled trial. *Arch. Phys. Med. Rehabil.* 100, 8 (Aug. 2019), 1400–1408.
  - [62] Shammugam Muruga Palaniappan and Bradley S Duerstock. 2018. Developing rehabilitation practices using virtual reality exergaming. In *2018 IEEE International Symposium on Signal Processing and Information Technology (ISSPIT)*. IEEE, 090–094.
  - [63] Cecily Partridge, Mathew Mackenzie, Susan Edwards, Andrea Reid, Sue Jayawardena, Nikki Guck, and Jonathon Potter. 2000. Is dosage of physiotherapy a critical factor in deciding patterns of recovery from stroke: a pragmatic randomized controlled trial. *Physiotherapy Research International* 5, 4 (2000), 230–240.
  - [64] Daniel Perez-Marcos, Odile Chevalley, Thomas Schmidlin, Gangadhar Garipelli, Andrea Serino, Philippe Vuadens, Tej Tadi, Olaf Blanke, and José d R Millán. 2017. Increasing upper limb training intensity in chronic stroke using embodied virtual reality: a pilot study. *Journal of neuroengineering and rehabilitation* 14 (2017), 1–14.
  - [65] Eva Pietrzak, Stephen Pullman, and Annabel McGuire. 2014. Using virtual reality and videogames for traumatic brain injury rehabilitation: a structured literature review. *GAMES FOR HEALTH: Research, Development, and Clinical Applications* 3, 4 (2014), 202–214.
  - [66] Wendy Powell, Vaughan Powell, and Maureen Simmonds. 2014. Virtual Reality for Gait Rehabilitation - Promises, Proofs and Preferences. In *Proceedings of the 7th International Conference on Pervasive Technologies Related to Assistive Environments* (Rhodes, Greece) (PETRA '14). Association for Computing Machinery, New York, NY, USA, Article 16, 7 pages. <https://doi.org/10.1145/2674396.2674450>
  - [67] Qinyin Qiu, Diego A Ramirez, Soha Saleh, Gerard G Fluet, Heta D Parikh, Donna Kelly, and Sergei V Adamovich. 2009. The New Jersey Institute of Technology Robot-Assisted Virtual Rehabilitation (NJIT-RAVR) system for children with cerebral palsy: a feasibility study. *J. Neuroeng. Rehabil.* 6, 1 (Nov. 2009), 40.
  - [68] Eric D Ragan, Doug A Bowman, Regis Kopper, Cheryl Stinson, Siroberto Scerbo, and Ryan P McMahan. 2015. Effects of field of view and visual complexity on virtual reality training effectiveness for a visual scanning task. *IEEE transactions on visualization and computer graphics* 21, 7 (2015), 794–807.
  - [69] Aimee P Reiss, Steven L Wolf, Elizabeth A Hammel, Erin L McLeod, and Erin A Williams. 2012. Constraint-induced movement therapy (CIMT): Current perspectives and future directions. *Stroke Res. Treat.* 2012 (April 2012), 159391.
  - [70] Marta Rodríguez-Hernández, Begoña Polonio-Moriano, Ana-Isabel Corregidor-Sánchez, José L Martín-Conty, Alicia Mohedano-Moriano, and Juan-José Criado-Álvarez. 2023. Can specific virtual reality combined with conventional rehabilitation improve poststroke hand motor function? A randomized clinical trial. *J. Neuroeng. Rehabil.* 20, 1 (April 2023), 38.
  - [71] Anca Salagean, Jacob Hadnett-Hunter, Daniel J Finnegan, Alexandra A De Sousa, and Michael J Proulx. 2022. A virtual reality application of the rubber hand illusion induced by ultrasonic mid-air haptic stimulation. *ACM Transactions on Applied Perception (TAP)* 19, 1 (2022), 1–19.
  - [72] Joseph P. Salisbury, Ted M. Aronson, and Tony J. Simon. 2020. At-Home Self-Administration of an Immersive Virtual Reality Therapeutic Game for Post-Stroke Upper Limb Rehabilitation. In *Extended Abstracts of the 2020 Annual Symposium on Computer-Human Interaction in Play* (Virtual Event, Canada) (CHI PLAY '20). Association for Computing Machinery, New York, NY, USA, 114–121. <https://doi.org/10.1145/3383668.3419935>
  - [73] Helle Spindler, Kasper Leerskov, Katrine Joensson, Gitte Nielsen, Jan Jesper Andreasen, and Birthe Dinesen. 2019. Conventional rehabilitation therapy versus telerehabilitation in cardiac patients: A comparison of motivation, psychological distress, and quality of life. *Int. J. Environ. Res. Public Health* 16, 3 (Feb. 2019), 512.
  - [74] Takashi Takebayashi, Kayoko Takahashi, Yuho Okita, Hironobu Kubo, Kenji Hachisuka, and Kazuhisa Domen. 2022. Impact of the robotic-assistance level on upper extremity function in stroke patients receiving adjunct robotic rehabilitation: sub-analysis of a randomized clinical trial. *Journal of NeuroEngineering and*

- Rehabilitation* 19, 1 (2022), 25.
- [75] Julia Tannus de Souza, Caroline Valentini, Eduardo Lazaro Martins Naves, and Edgard Afonso Lamounier Jr. 2022. A Virtual Reality Exergame with a Low-Cost 3D Motion Tracking for At-Home Post-Stroke Rehabilitation. In *Symposium on Virtual and Augmented Reality* (Virtual Event, Brazil) (SVR'21). Association for Computing Machinery, New York, NY, USA, 172–176. <https://doi.org/10.1145/3488162.3488223>
  - [76] Kelly O Thielbar, Thomas J Lord, Heidi C Fischer, Emily C Lazzaro, Kristin C Barth, Mary E Stoykov, Kristen M Triandafilou, and Derek G Kamper. 2014. Training finger individuation with a mechatronic-virtual reality system leads to improved fine motor control post-stroke. *J. Neuroeng. Rehabil.* 11 (Dec. 2014), 171.
  - [77] Maferima Touré-Tillery and Ayelet Fishbach. 2014. How to measure motivation: A guide for the experimental social psychologist. *Social and Personality Psychology Compass* 8, 7 (2014), 328–341.
  - [78] KI Ustinova, J Perkins, WA Leonard, and CJ Hausbeck. 2014. Virtual reality game-based therapy for treatment of postural and co-ordination abnormalities secondary to TBI: a pilot study. *Brain injury* 28, 4 (2014), 486–495.
  - [79] Giulio Verrienti, Cecilia Raccagni, Ginevra Lombardozzi, Daniela De Bartolo, and Marco Iosa. 2023. Motivation as a measurable outcome in stroke rehabilitation: A systematic review of the literature. *Int. J. Environ. Res. Public Health* 20, 5 (Feb. 2023).
  - [80] Nicolas Wenk, Mirjam V Jordi, Karin A Buetler, and Laura Marchal-Crespo. 2022. Hiding assistive robots during training in immersive vr does not affect users' motivation, presence, embodiment, performance, nor visual attention. *IEEE transactions on neural systems and rehabilitation engineering* 30 (2022), 390–399.
  - [81] Carla Winter, Florian Kern, Dominik Gall, Marc Erich Latoschik, Paul Pauli, and Ivo Käthner. 2021. Immersive virtual reality during gait rehabilitation increases walking speed and motivation: a usability evaluation with healthy participants and patients with multiple sclerosis and stroke. *J. Neuroeng. Rehabil.* 18, 1 (April 2021), 68.
  - [82] Shih-Ching Yeh, Si-Huei Lee, Rai-Chi Chan, Yi Wu, Li-Rong Zheng, and Sheryl Flynn. 2017. The efficacy of a haptic-enhanced virtual reality system for precision grasp acquisition in stroke rehabilitation. *J. Healthc. Eng.* 2017 (Nov. 2017), 9840273.
  - [83] Francesco Zanatta, Naima Z Farhane-Medina, Roberta Adorni, Patrizia Steca, Anna Giardini, Marco D'Addario, and Antonia Pierobon. 2023. Combining robot-assisted therapy with virtual reality or using it alone? A systematic review on health-related quality of life in neurological patients. *Health Qual. Life Outcomes* 21, 1 (Feb. 2023), 18.
  - [84] André Zenner and Antonio Krüger. 2019. Estimating detection thresholds for desktop-scale hand redirection in virtual reality. In *2019 IEEE Conference on Virtual Reality and 3D User Interfaces (VR)*. IEEE, 47–55.
  - [85] Weiwei Zhao, Chuanguang Ju, Daozhen Wang, and Huifen Shen. 2019. Clinical observation of effects of ultrashort wave therapy combined with acupuncture and rehabilitation training in the treatment of patients with dysphagia after stroke. *Journal of Neurorestoratology* 7, 3 (2019), 136–142.
  - [86] Lukas Zimmerli, Mario Jacky, Lars Lünenburger, Robert Riener, and Marc Bolliger. 2013. Increasing patient engagement during virtual reality-based motor rehabilitation. *Arch. Phys. Med. Rehabil.* 94, 9 (Sept. 2013), 1737–1746.